International Journal of Recent Advances in Multidisciplinary Topics
Volume 7, Issue 6, June 2026
www.ijramt.com | E-ISSN: 2582-7839 | RESAIM Publishing | www.resaim.com

Analysis of the Increased Tenderness for Alcohol Amongst Women and Factors Influencing It

Naiya Rana Ratnoday[1], Nilesh Ratnoday[1*]
[1]*Assistant Professor, Institute of Hotel Management, Guru Kashi University, Talwandi, India*

***Abstract*: Background: Alcohol consumption among women has steadily risen globally, raising concerns about its potential health and social consequences. This study investigates different factors that influence this, some of them are fun, stress, life events, and many more. The study aims to devote research attention to women's drinking factors, their judgement, their knowledge about ancient women consuming alcohol, health problems they are aware of, and if they feel that there is a gender specific difference in consuming alcohol. Methodology: The research will employ a mixed-methods approach, combining quantitative and qualitative data collection. Conclusion: About this research, some women do not have much awareness about the health problems, but still, they tend to consume more alcohol, and also women do not have any knowledge about the ancient alcohol consumption by women, and today also women face societal judgement because of their drinking habits. The increase in consumption is seen in teenagers and Generation Z, including students. Professionals from the IT sector was found to be more inclined towards alcohol. Most women are aware of the health effects. The factors to consume alcohol tend to be many, but fun and social gatherings are the most common.**

***Keywords*: Alcohol, Consumption, Health Problems.**

1. Introduction

A. Historical Backdrop of Indian Alcohol

Ancient India Had a Rich History of Alcohol production and consumption. The peculiarity of liquor utilization has happened somewhat all through written history; researcher have found proof of wine and lager in earthenware containers at the locales of a Sumerian general store in Western Iraq that goes back 6,000 years. Alcohol came to India way earlier that anyone would guess. Indians were the first civilizations to discover the science of distillation (Fairley, 1906) Inside the Indian setting, inebriating drinks were known even in the Vedic and post Vedic times and the Apparatus Veda has one section that goes on about soma which is accepted to have been the heavenly beverage during the Vedic time frame. Indeed, even in the legendary Mahabharata reference has been made to alcohol which appears to have been utilized to obliterate Yadavas. The principal notice of liquor shows up just with the Vedas, when the Rigveda (1700 BC) discusses the inebriating like soma and Praha mana. Hymns Praising Soma have been found through the ages & 114 hymn entirely is devoted to the purification of soma. (Singh, N. L., Mishra, P. K., Shukla, S. K., Kumar, J., & Singh, R. (2010).)

The most normally realized old Indian alcohol was Soma, referenced in Hindu sacred texts like the Rigveda. Soma was a custom beverage utilized in strict functions and was accepted to have psychedelic properties. One more well-known liquor in antiquated India was Madya, produced using matured rice, natural products or blossoms. Madya was consumed by individuals from varying backgrounds, including eminence, and was additionally utilized in friendly and strict occasions. Moreover, there were territorial cocktails like Sura produced using rice or grain, and Drink produced using palm sap. These refreshments were delivered and consumed locally and assumed a significant part in the economy and day to day existence of the networks that created them.

The Brahmins, the yogic, accepted that soma furnished them with a something beyond a hotline to the heavenly; it pervaded them with extraordinary powers. (Jadavji Tricumji Acharya & Narayan Ram Acharya(ed.),1987)

Research on drinking designs among ladies by and large shows that ladies drink not as much as men (Wilsnack, 1996), Truth be told, in all nations and verifiable periods examined, men have polished off additional liquor and revealed more liquor related issues than ladies (Fillmore et al., 1997).

In the territories of Gujarat, Nagaland, Mizoram and Manipur, there stays authorized restriction, which has driven up the interest for unlawful liquor (Malhotra & Mohan, 2000). The risk of drinking alcohol is rising along with India's growing passion for it. Various factors make it important to pay attention when female alcohol consumption rises. According to estimates, more than 140,000 persons per year die from alcohol-related causes (about 97,000 males and 43,000 women). (Alcohol and Public Health: Alcohol-Related Disease Impact. 2016)

For some time, it has been reported that female alcohol use is progressively increasing, especially among younger age groups of women. (Australian Institute of Health and Welfare. 1998)(Wilsnack SC, Wilsnack RW, Klassen AD. 1983±84, 3 ±13.) Research indicates that women are using and abusing alcohol at higher rates. (White A, Castle IJP, Chen CM, Shirley M, Roach D, Hingson R. 2002 to 2012) (Slade T, Chapman C, Swift W, Keyes K, Tonks Z, Teesson M. 2016) The creation volume of ethanol in India for 2020 was projected to associate with three billion Liters, higher than the ethanol creation

*Corresponding author: nratnoday@gmail.com

volume in 2019. This ethanol was utilized in refreshments, fuel and other modern synthetic substances in the country. (Madhumitha Jaganmohan, 2022)

B. Alcohol Consumption Among Ancient Indian Women

XI: 13. Drinking (spirituous alcohol), partner with underhanded individuals, partition from the spouse, meandering aimlessly abroad, sleeping (at untimely hours), and abiding in other men's homes, are the six reasons for the destruction of ladies.

XI: 80. A spouse who drinks spirituous alcohol, is of terrible direct, defiant, infected, naughty, or inefficient, may out of the blue be supplanted (by another wife). – The Laws of Manu c. 200 A.D.

Liquor was regularly utilized in labour all through the ages and lager was remembered to strengthen a person for bosom taking care of (Galactagogue). Hot drinks of different natural product tonics have been utilized for quite a long time for feminine spasms and for pre-feminine strain. Liquor was accepted to have restorative properties and various creations which included liquor were given to small kids the two guys and females in Goan culture. Ladies that drank openly or became smashed were viewed as physically rash and flippant moms. Despite the fact that liquor is effectively accessible, served at parties, utilized with respect to restorative purposes and for pressure decrease, ladies were supposed to keep their toasting negligible socially satisfactory levels and not become inebriated (Hanson, J., 2006).

C. Alcohol Consumption Among Modern Women

All through Western history, alcoholic ladies have been exposed to additional limitations than alcoholic men and have been punished all the more brutally for challenging codes against drinking (Sandmaier, 1992). Liquor addiction was once viewed as an issue for men, yet an assessed 3.9 million, or around 33% of all liquor manhandling or liquor subordinate people, are ladies (NIAAA, 1995).

A large portion of ladies would drink periodically or at social capabilities and were well inside the constraints of generally safe utilization until the event of intense life stressors (models like demise of spouse quite early on, death of a girl through a mishap and so forth). To numb the agony and to have the option to rest these ladies had their most memorable beverage to adapt to harming. Not understanding the cyclic impacts of fixation these ladies wound up in a winding moving downwards towards a far more regrettable condition than at the beginning of the emergency circumstance. Their aggravation was misjudged and that deteriorate what is going on till pressing forward was the only option. They were peered downward on 'as though they were hoodlums 'or 'with free ethics'.Double standards of drinking that judge unnecessary (or any) liquor utilize more cruelly for ladies than for men have been accounted for in many societies since the beginning of time (e.g., Blume, 1997). In such society, these ladies with drinking issues, find it hard to look for help and once in a while are kept by embarrassed relatives from looking for treatment. Across the globe, endeavours are being had to concentrate on this effect as reports are showing that the hole is gradually shutting particularly with youth Women consumed wine in the greatest amount, whereas men chiefly consumed beer (Mukamal, K. J., Tolstrup, J. S., Friberg, J., Jensen, G., & Grønbæk, M. (2005).

The pervasiveness of drinking, as well as AUD (alcohol use Disorder), among more established ladies, is extremely low and is reliable with the overall epidemiological picture in India (Murthy, P., Manjunatha, N., Subodh, B.N., Chand, P.K., & Benegal, V. (2010)). Drinking among more established ladies is considerably less continuous than among men in ongoing examinations (Alam and Karan,2011). It should be noted, nonetheless, that liquor utilization patterns among grown-up ladies in India is changing rap inactively (Ray S, 2002)There is likewise an inescapable idea that liquor use is restricted to ancestral ladies, ladies of lower financial status, business sex labourers and to a restricted elite of the rich, and not leaned toward by ladies from the centre or upper financial classes (Ray and Sharma, Kumar & Parthasarathy, 1994).Ladies drank as much as men when they drank (Benegal, V., Gururaj, G., & Murthy, P., 2003).Misogynist perspectives to ladies' drinking continue everywhere (Blume, 1997), and in many societies there is a more prominent disgrace connected to drinking by females than among guys.

An illustration of this is a directive contained in the old Brahminical text on private regulation, the Manu smriti (around 200 A.D.), which cautions that "drinking (spirituous alcohol), partner with mischievous individuals, partition from the spouse, meandering abroad, resting (at unexpected hours), and abiding in other men's homes, are the six reasons for the destruction of ladies" (Doniger,1991).Ultimately nonetheless, ladies' utilization of liquor was confined to culinary and restorative purposes, albeit high society women were allowed to Savor liquor modest quantities at their homes or confidential get-togethers. In Goa, in light of the impact of the Portuguese traditions and customs, ladies would likewise drink limited quantities at parties. As a matter of fact, at functions like weddings, birthday celebrations or commemorations gatherings, the raising of the toast is gone with a standard glass of the best wine, regularly champagne. The presentation of new items, for example, seasoned and gentle alcoholic items is planned to enrol non-drinkers and is designated fundamentally at ladies" (Naik, 2002).

Up to this point, the mentality that 'women don't drink' has generally kept female drinking and liquor maltreatment from public conversation. This disposition, be that as it may, has been proposed rather as of late for example throughout the previous 100 years or somewhere in the vicinity, preceding which liquor and ladies have been chronicled to be laced since forever ago. In a few old societies, ladies had a functioning part in the drinking services and fables, including directing at the Greco-Roman factions of Dionysius and Bacchus and fermenting brew in old Babylon as sanctuary priestesses. The Egyptians considered the information on the most proficient method to blend brew as a gift bestowed to them by their goddess of nature, Isis (Hornik, 1977).

Accordingly, the number and effect of more established female consumers is supposed to increment throughout the

following 20 years as the difference among people's drinking rates decline. Because of contrasts in digestion of liquor, ladies of any age contrasted with men are at higher gamble for negative physical, clinical, social, and mental outcomes related with in danger and more significant levels of liquor utilization. Maturing ladies face new arrangements of precursors connected with difficulties in the centre and more established grown-up periods of life, like menopause, retirement, "void home," restricted versatility, and sickness. (Epstein, E. E., Fischer-Elber, K., & Al-Otaiba, Z. (2007)

Enormous Liquor has taken advantage of the Coronavirus pandemic in India to change ladies over completely to liquor utilization. By far most of ladies in India live liquor free. The World Wellbeing Association reports that in 2016 just about 3/4 of the female populace in India north of 15 years old lived liquor free (74.6% beyond a year liquor teetotaller; 68.8% lifetime liquor teetotallers). Distressing occasions correspond with more elevated levels of liquor utilization at the populace level," said Ruler Singhal, Pioneer, of Local area Against Intoxicated Driving (CADD), according to The New Indian Express.

D. Reason of Increase in Consumption

Rising affluence, aspirations, societal pressure and exposure to a different lifestyle is driving women to experiment with alcohol, stated the survey, conducted by the Community Against Drunken Driving (CADD) among 5,000 women aged between 18 to 70 in Delhi. Liquor utilization among ladies has in the previous many years started to draw in expanded consideration. This seems to have come about because of the ascent of woman's rights, expanded drinking by ladies, and wellbeing concerns, for example, 'hidden alcoholism' ' and cirrhosis of the liver (Chalfant and Roper, 1980). Naturally, no country in this world is liberated from the liquor threat. Be that as it may, there is little information accessible on the degree of liquor abuse or liquor maltreatment in India and all the more explicitly its connection to ladies' wellbeing and since drinking is presently not an untouchable for working class metropolitan Indian ladies; reports of Ladies' liquor utilization are of arising concern. (Ravindran.S.,2010)

Changing patterns of female alcohol consumption reflect societal changes whereby it has become more acceptable for women to consume alcohol, and to do so in ways that mirror drinking patterns of their male counterparts ([Plant M. Women and alcohol: contemporary and historical perspectives. 1997.)

Drinking alcohol as a means to socialize and battle personal problems. (survey conducted by CADD) An investigation of drinking propensities led in Bangalore city revealed that a fifth of youngsters who regularly visited bars on ends of the week were young ladies matured between 13-19 years (Kumar, 1997) In the worldwide writing, changes in ladies' schooling, work, societal position and monetary freedom have been ensnared in the assembly of male and female drinking designs. (Neve, R. J. M., Lemmens, P. H., and Drop, M. J. (1996).) held that as ladies slowly obtain equivalent freedoms in work and social circumstances, this could likewise be reflected in a penchant to drink in equivalent sums and in similar way as men. While this may not make sense of changes in that frame of mind of non-working females, it mirrors a general change in cultural mentalities to ladies' drinking.

E. Comparison of Male and Female Alcoholic Impact

There is research that concentrates on showing that ladies begin to have liquor related issues sooner and at lower drinking sums than men and in light of several factors. (Erol A, Karpyak 2015) Overall, ladies weigh not as much as men. Likewise, liquor dwells prevalently in body water, and pound for pound, ladies have less water in their bodies than men. (Kalant H. 1971.). This truly intends that after a lady and a man of a similar weight drink a similar measure of liquor, the lady's blood liquor focus (how much liquor in the blood) will quite often be higher, putting her at more serious gamble for hurt. For instance, research proposes that ladies are more probable than men to encounter headaches and liquor actuated power outages at the amount portions of liquor. (Vatsalya V, Stangl BL, Schmidt VY, Ramchandani, 2018) (Hingson R, Zha W, Simons-Morton B, and White A. 2016) Ladies, contrasted with men, have had bigger expansions in liquor related crisis office visits, hospitalizations, and passing's in the beyond 20 years. (White AM. 2020;40(2))

Moreover, ladies' general gamble taking way of behaving, (for example, drink driving and cooperation in crime is likewise expanding at a raised rate in correlation with men. (Australian Institute of Health and Welfare. 1998)[Polk K.2001.) Appropriately, ladies' liquor utilization in regards of its examples, antecedents and consequences is unique in relation to the liquor utilization of men.

In overall communities concentrates all through the world, when contrasted with ladies, men are all the more frequently consumers, drink more liquor, and create more issues thusly (Almeida et al., 2004; Fillmore et al., 1991; Hao et al., 2004; Jhingan et al., 2003; Kebede and Alem, 1999; Yamamoto et al., 1993). The size of these distinctions in sexual orientation fluctuates extraordinarily from one society to the next. The orientation hole in drinking conduct is one of only a handful of exceptional general distinctions in sexual orientation in human social way of behaving that actually wins. These distinctions in sexual orientation in liquor use have supported expensive predispositions in how social orders recognize and attempt to control liquor related issues. From one viewpoint, the relationship of weighty drinking with showcases of manliness or male fellowship might urge male consumers to deny or limit issues or dangers coming about because of their drinking, or to see tanked conduct as typical or reasonable, in any event, when it prompts savagery. Then again, suspicions that ladies don't drink vigorously may at first lead to ladies' drinking issues being limited or disregarded (for instance, by clinical specialists; (Brienza and Stein, 2002; Svikis and Reid-Quinones, 2003; Weisner and Matzger, 2003), yet when ladies' liquor misuse or reliance becomes prominent, the social response might move from aloofness to shock and endeavours to rebuff ladies who drink in socially disliked ways. (Abel and Kruger, 2002; Blume, 1997; De Ville and Kopelman, 1998)

F. Alcohol Impact on Health

Worldwide examination has laid out that liquor utilization is a gamble factor for in excess of 200 illness and injury conditions and that 5.1% of the worldwide sickness trouble has been credited to hurtful liquor utilization. WHO assesses that destructive utilization of liquor brings about roughly 3.3 million passings every year, or 5.9% of all passings. (WHO,2018) Past its wellbeing results, the unsafe utilization of liquor causes huge financial effects for people, families, and society at large. these effects range from lost efficiency in the working environment, to joblessness, crimes and socially degenerate way of behaving, property misfortune and harm, as well as profound pain and psychosocial influence on family and local area individuals.

Across India utilization of liquor is answerable for almost 1.4 million instances of 19 diseases for which, liquor has been distinguished as an unmistakable gamble component and 11 million inability changed life years (DALYs) which is 8.8% of the all-out DALYs each year. It is assessed that liquor utilization brings about almost 2.4 million instances of violations enrolled with the police, 1.6 million cases brought to preliminary. (WHO, 2018) The male: female sex proportion for head and neck disease has dropped quickly in the UK from around 5 :, a long time back to 2 : 1 as of now. The frequency has ascended from 5.14 per 100 000 out of 1989 to 5.71 in 1998 in ladies while remaining consistent in men. Liquor utilization in ladies during the same ten years has similarly shown an increment. (Viswanathan, H., & Wilson, J. A. (2004). Where ladies and men drink at similar rate, ladies keep on being at higher gamble than people for specific serious clinical results of liquor use, including liver, mind, and heart harm. In spite of the fact that we live in a general public that Favors fairness between the genders, people are not equivalent when it includes being impacted by liquor (Edith.S., 1986.; Frezza, M.;Taylor, J.L.; Dolhert, N.; Friedman, L.; et al. 1996; NIAAA, 2006). Overall, ladies who drink polish off less liquor and have less liquor related issues and reliance side effects than men. (Malin, H., Coakley, J., and Kaelber, C., 1982), but among the heaviest consumers, ladies equivalent or outperform men in the quantity of issues that outcome from their drinking (Wilsnack, R.W., Wilsnack, S.C., and Klassen, A.D., 1984). Liquor use and manhandle likewise add to wounds, car accidents, and brutality. Liquor can uniquely influence labourer efficiency and non-appearance, family associations, and school execution, and it can kill, straightforwardly through mishaps or in a roundabout way through organs disappointment or sickness. Liquor consequently is a serious general medical condition. (Room, Babor, and Rehm, 2005)

Even though all alcohol abuse has substantial public health risks, women who drink are more likely than men to experience specific alcohol-related issues. (· Erol A, Karpyak V. 2015,1-13) The more noteworthy concern anyway is additionally that the age gathering of respondents for starting drinking is lessening and even young ladies, more youthful than the lawful age of 18 years are hard-core boozing. India is showing a remarkable expansion in liquor utilization with the commencement age on a disturbing lessening. Customarily the distinctions in sexual orientation demonstrated that the quantity of ladies drinking liquor was low, however powerful showcasing and propelling way of life can roll out a critical improvement in this proportion which is becoming obvious across India. (Benegal V., 2003)

2. Literature Review

Barik, A., Rai, R. K., & Chowdhury, A. (2016). Alcohol use-related problems among a rural Indian population of West Bengal: an application of the Alcohol Use Disorders Identification Test (AUDIT). *Alcohol and alcoholism*, *51*(2), 215-223 founds that more than 19% of guys and 2.4% of females were current liquor clients. Mean ethanol utilization on a regular drinking day among guys was assessed to be higher (96.3 gm) than females (56.5 gm). Mean Review score was 11 among current liquor clients. Review displayed in the arranged logit relapse assessed liquor use-related issues to be low among ladies, Booked Clans and unmarried individuals, though liquor use-related issues enlisted high among Muslims. the end from this paper was provincial populace gives off an impression of being needing a viable intercession program,

Benegal, V., Nayak, M., Murthy, P., Chandra, P., & Gururaj, G. (2005). Women and alcohol use in India. *Alcohol, gender and drinking problems: Perspectives from low- and middle-income countries*- mentioned that 5.8 % of every single female respondent (86 out of 1471) detailed drinking liquor somewhere around once over the most recent a year, contrasted with liquor use in 33.2 % of every male respondent (502 out of 1508). 56% of the men and 84% of the ladies were lifetime teetotallers, while 10.5 % of men and 10.1% ladies detailed that they were current teetotallers with a background marked by liquor use (something like once) in the course of their life.

Chowdhury, A. N., Ramakrishna, J., Chakraborty, A. K., & Weiss, M. G. (2006). Cultural context and impact of alcohol use in the Sundarban Delta, West Bengal, India. *Social Science & Medicine*, *63*(3), 722-731. More than 19% of guys and 2.4% of females were current liquor clients. Mean ethanol utilization on a regular drinking day among guys was assessed to be higher (96.3 gm) than females (56.5 gm). Mean Review score was 11 among current liquor clients. Review displayed in the arranged logit relapse assessed liquor use-related issues to be low among ladies, Booked Clans and unmarried individuals, though liquor use-related issues enlisted high among Muslims. the end from this paper was provincial populace gives off an impression of being needing a viable intercession program, maybe focusing on men and the family, pointed toward decreasing the degree of liquor use and related issues.

Das, S. K., Balakrishnan, V., & Vasudevan, D. M. (2006). Alcohol: Its health and social impact in India. *National Medical Journal of India*, *19*(2), 94. Mentioned that the example of liquor admission all over the planet is continually advancing. There is in this way a solid defence for the wellbeing calling to move forward its wellbeing backing as for strategies to lessen paces of liquor utilization. Despite the fact that India is by and large viewed as a dry nation, drinking rehearses vary impressively among individuals living in the southern and northern regions, and even among individuals from various positions dwelling in a similar locale. Because of the social and

ethnic varieties in drinking rehearses, no single meaning of 'issue drinking' or liquor reliance exists that can be consistently applied in all nations or societies. Thusly, scientists, clinicians and general wellbeing authorities at enticing to foster compelling anticipation and treatment approaches should consider the populace's perspectives and assumptions about liquor utilization and its belongings.

Dawson, D. A., & Archer, L. (1992). Gender differences in alcohol consumption: effects of measurement. *British journal of addiction*, *87*(1), 119-123. Mentioned that the Information from a 1988 review of US drinking propensities and related issues uncovered contrasts in male and female examples of liquor utilization. Men were more probable than ladies to be current consumers (64 v. 41%), and lager represented a bigger extent of their general admission. Men's typical everyday ethanol admission was about two times just that high of ladies, 17.5 v. 8.9 g. Change for contrasts in body weight and arrangement considerably diminished the male-to-female proportion of utilization. Men were more probable than ladies to be named weighty consumers, and the abundance extent of guys so arranged expanded with the seriousness of the proportion of weighty drinking.

Gururaj, G., Gautham, M. S., & Arvind, B. A. (2021). Alcohol consumption in India: A rising burden and a fractured response. *Drug and Alcohol Review*, *40*(3), 368-384. states that the predominant liquor control approaches and projects in India have been not exactly completely successful in controlling the weight of liquor use and its related effect. There is a requirement for complete, proof based and agreement driven public liquor control strategy to guide and support the Indian states in managing liquor and diminishing the related weight fittingly. Compelling executions of such approaches are vital to its prosperity.

Jha, P., Ranson, K., Nguyen, S.N., Yach, D., 2002. Estimates of global and regional smoking prevalence in 1995, by age and sex. Am. J. Public Health 92, 1002–1006. The more noteworthy recurrence of tobacco use in Indian men versus ladies is predictable with 1995 appraisals of generally speaking smoking pervasiveness for men of 40% and 4% for ladies in the South Asia locale.

Jonas, J. B., Nangia, V., Rietschel, M., Paul, T., Behere, P., & Panda-Jonas, S. (2014). Prevalence of depression, suicidal ideation, alcohol intake and nicotine consumption in rural Central India. The Central India Eye and Medical Study. *PLoS One*, *9*(11), e113550. Gentle to direct sorrow (Community for Epidemiologic Investigations Wretchedness Scale CESD aggregate score: 15-21) was recognized in 1862 (39.6%) people (33.5% of men, 44.8 of ladies), and significant sorrow (CESD total score .21) in 613 (13.0%) people (8.1 of men, 17.3% of ladies). Self-destruction endeavour was accounted for by 199 (4.2%) members and self-destructive contemplations during the most recent a half year by 238 (5.1%) people. Liquor utilization was revealed by 1081 (23.0%) members; 283 (6.0%) subjects had a Review score $8 (risky drinking), and 108 (4.63%) subjects a score $13 (ladies) or $15 (men) (liquor reliance) In provincial Focal India, predominance of significant sadness was tantamount to figures announced from other agricultural nations. Predominance of smoking and dangerous liquor utilization was higher than as announced from metropolitan districts. Measures ought to be taken to address the generally high predominance of self-destruction endeavours and considerations on self-destruction in provincial Focal India.

Kirmani, M. N., & Suman, L. N. (2010). Out of these, 150 (79%) were boys and 40 (21%) were girls. Beer was the most frequently consumed alcohol beverage by both boys and girls. An almost equal number of boys (23%) and girls (26%) reported alcohol use in their families. However, a greater number of boys (32%) reported that alcohol use was prevalent among their peers compared to girls (11%).

Kumar, K., Kumar, S., & Singh, A. K. (2018). The Prevalence and socio-demographic correlates of alcohol consumption: Survey findings of 5 different states Drug *and alcohol dependence*-mentioned that more than 33% of the example respondents (38.6%, 95%CI = 29.2-48.8%) answered to be current consumers and around one-fifth (21.7%, 95%CI = 4.2-31.7%) were weighty consumers and 7.4% (95%CI = 4.6-11.6%) were weighty verbose consumers.

Office for National Statistics, London mentioned the 1995 changed benchmarks for reasonable drinking (3-4 units/day for men and 2-3 for ladies), expects that it won't build huge wellbeing risk however ongoing examinations have connected expanded risk specifically of bosom malignant growth in ladies with lower levels of liquor utilization. In Britain in 2000, 81% of individuals matured 16 and over had heard of estimating liquor utilization in units, yet over 45% of respondents who said that they had known about day-to-day benchmarks didn't have the foggiest idea what they were.

Luca, D. L., Owens, E., & Sharma, G. (2019). The effectiveness and effects of alcohol regulation: evidence from India. *IZA Journal of Development and Migration*, *9*(1), 1-26. Stated that considerable proof that lessening admittance to liquor through MLDA regulations diminishes the probability of drinking, smoking, and aggressive behaviour at home.

A. Family, Parental Impacts, Youngster Raising Practices

Various investigations have discovered that juvenile drinking was related with drinking by relatives (e.g., Goodwin, 1985; Hansen et al., 1987; Rachal et al., 1982). Guardians who permitted their 5th grade kids to drink at home were bound to have youngsters who drank liquor later (Jackson et al., 1999). Additionally, offspring of guardians who drank were less inclined to see drinking as destructive (Hawkins et al., 1997). Despite the fact that coinciding among juvenile and family drinking might be the consequence of either hereditary or ecological elements, research directed with twins and different kin found a social virus aspect to young adult drinking that was free of hereditary relatedness (Rende et al., 2005), proposing that social impacts are essential variables in starting drinking. As indicated by Mcord (1965) Substance use has been credited to family issues especially early isolating from one or both the guardians. Mowrer (1942) sees the family as focal in the production of the character yet considers dependence on be the consequence of various humanistic and mental reasons. From (1941) propose weakness, and a need to submit to another

authority as reason for compulsion and considers addiction on be a sadomasochist quality in the addict.Chein (1969) joins addiction on the level of family profound wellbeing. Mohan d. et al (1980) viewed that as a bigger number of users than non-users are disappointed with 15.83% and 8.63 % or in various 29.9% and 23.7% to their current family life. Fergusson et al (1979) observed that there is connection between parts of family life and differential contribution in habit-forming drugs and liquor. Consequences of studies highlighted a pessimistic relationship between level of addiction and emotional sensation of fulfilment and inclusion with as well as the capacity to hand-off upon the family as a unit. Lewis and Osberg (1958) additionally noticed that the family addiction showed absence of cohesiveness. Anyway Rosenberg brought up that not exactly 50% of his subject addicts came from family foundation where the addicts had the two parents live with them basically till the age of 15 yrs.Roseberg (1971), Chein and Blum (1972) recommend that the presence or nonappearance of parent is less essential than the idea of the parent youngster relationship. In the event that the parental relationship is positive, the youngster is less inclined to become associated with drugs. The simple presence of a parent appears to be irrelevant on the off chance that that parent isn't successfully engaged with the youngster.

B. Social Factors

Windham and Aldridge (1965) as well as Siegel (1952) referred to the conventional conviction that the utilization and maltreatment of liquor was a male right.

Clark (1967) proposed that drinking distinctions depend to a great extent on the assumption that female sex jobs are described by what he named 'customariness'. By this he referred to the 'acknowledgment of the predominant 'official' principles of ethical quality and legitimacy'.

Preston (1964) comparatively stated that drinking is many times an image that separates the genders. Windham and Aldridge (1965) moreover brought up that ladies customarily would in general be financially subject to and subordinate to men.

In such manner Knupfer et al. (1963) stressed that "different individuals from the populace who don't make money by paid work likewise have more limited drinking honours than grown-up free guys for instance, kids, detainees, mental patients, and people on help".

Wilsnack and Wilsnack (1978) bring up that expanded drinking among females may be a consequence of the ladies' development and changes in ladies' jobs, particularly changes that include openness to previously manly conditions and jobs. They recommend that adjustments of sex jobs could expand ladies' openness to liquor and chances to drink; could adjust customary standards against female drinking, accordingly making drinking more passable; furthermore, could offer females new objectives and yearnings, subsequently causing pressure that liquor may be utilized to decrease.

Mishra, V. K., Srivastava, S., & Murthy, P. V. (2021) mentioned that Adult Indian women's population-attributable risk for multimorbidity: Do smoking tobacco, biting tobacco and polishing off liquor have an effect *PloS one*, *16*(11), e0259578. Mentioned that Around 1.6% of ladies had multimorbidity in India. The pervasiveness of multimorbidity was high among ladies from southern locale of India. Ladies who smoke tobacco, bite tobacco and drink liquor had 87% [AOR: 1.87CI: 1.65, 2.10], 18% [AOR: 1.18; CI: 1.10, 1.26] and 18% [AOR: 1.18; CI: 1.04, 1.33] altogether higher probability to experience the ill effects of multi-horribleness than their partners individually. Populace Inferable Gamble for ladies who smoke tobacco was 1.2% ($p<0.001$), bite tobacco was 0.2% ($p<0.001$) and it was 0.2% ($p<0.001$) among ladies who polished off liquor his examinations show the significant job of way of life and conduct factors like smoking and biting tobacco and drinking liquor in the commonness of multimorbidity among grown-up Indian ladies. The subgroups distinguished as at expanded risk in the current review can be focused on while going with strategies and wellbeing choices and suitable comorbidity the board can be carried out.

Moinuddin, A., Goel, A., Saini, S., Bajpai, A., & Misra, R. (2016) reasoned that general liquor utilization all through the world is on an ascent with a reliable restricting orientation hole in spite of the fact that disparity is available inside countries and social orders.

Mondal, N. A., Mog, M., & Khan, K. A. (2018) stated the liquor utilization was two percent higher in 2004-05 (37%) than the new review of IHDS (2011-12) (35%) in India. Besides, there were a couple of states where the usage of alcohol had extended from 2004 to 2012 like Mizoram (44%), Kerala (19%), and Jammu and Kashmir (16%). In opposition to that, there were additionally a couple of states where liquor utilization had diminished from 2004 to 2012 like Rajasthan (8%) and Tamil Nadu (5%).

Murthy, P. (2015) said that social settings of liquor use can't be separated from socio-political settings. In India, religion was major areas of strength for an of restraint and social request was areas of strength for an of purpose and overabundance

Nadkarni, A., Murthy, P., Crome, I. B., & Rao, R. (2013). Alcohol use and alcohol-use disorders among older adults in India: a literature review. *Aging & mental health*- mentioned that more established ladies are for the most part liable to be liquor abstainer. The general pattern has all the earmarks of being a decrease in current drinking among progressive age companions beyond 50 years old.

O'Malley, S. S., Sinha, R., Grilo, C. M., Capone, C., Farren, C. K., McKee, S. A., & Wu, R. (2007) mentioned No tremendous contrasts were seen on the essential results of time to first drinking day, time to first day of weighty drinking, or the level of members who kept on gathering the standards for liquor reliance. Optional examinations uncovered that naltrexone fundamentally deferred the chance to the second (w2 5 5.37, p 5 0.02) and third (w2 5 4.35, p 5 0.04) drinking days among subjects who didn't keep up with forbearance from liquor. Among those with eating unsettling influences, side effects of eating pathology improved during treatment, however the impacts didn't vary as indicated by drug condition. At the point when utilized related to h Mental Conduct Adapting Abilities Treatment (CBCST), naltrexone didn't fundamentally further develop savouring results the general example of liquor

subordinate ladies. Notwithstanding, naltrexone might be good for ladies who can't keep up with complete restraint from liquor. For ladies with simultaneous eating pathology, cooperation in treatment for liquor abuse might be related with enhancements in eating pathology.

Pfefferbaum, A., Rosenbloom, M., Deshmukh, A., & Sullivan, E. V. (2001). Sex differences in the effects of alcohol on brain structure. *American Journal of Psychiatry*, *158*(2), 188-197. Ladies, paying little heed to finding, had less cortical dark and white matter and more modest third ventricles than men, reliable with sex-related contrasts in intracranial volume. Drunkards had bigger volumes of cortical sulci and sidelong and third ventricles than examination subjects.

Rahman, L. 2003. Alcohol prohibition and addictive consumption in India. Leverhulme Centre for Research on Globalization and Economic Policy (GEP), University of Nottingham shows that liquor forbiddance strategies in India are related with a 20-40% decline in the likelihood of liquor utilization and a 40% reduction in the amount of liquor consumed utilizing family consumption information.

Rastogi, A., Manthey, J., Wiemker, V., & Probst, C. (2022). mentioned that (Life-time Abstinence) LA, AUD (Alcohol use Disorder) and QU (current drinkers) information were accessible for 31, 36 and 12 states/UTs, separately. Cd model {(current drinking (CD) (12 months)} appraisals went from 6.4% (95% CI = 2.1%-18.1%; guys) in Lakshadweep and 1.3% (95% CI = 0.7%-2.6%; females) in Delhi to 76.1% (95% CI = 68.1%-82.6%; guys) and 63.7% (95% CI = 49.4%-75.7%; females) in Arunachal Pradesh. Over the long run, Compact disc diminished in many states/UTs in the noticed information, going against expanding public level APC (alcohol per capita consumption) patterns. Liquor use (estimated as predictable current drinking) in India has huge provincial varieties, with liquor utilization being most pervasive in the North-East, Chhattisgarh, Telangana, Himachal Pradesh, Punjab and Jharkhand.

At the point when this happens the mind is probably going to support more long-lasting harm, experienced as memory hindrance (NIAAA, 2004).

Speculative Models of Etiology. (field concerned with the study of causes, origins, or reasons behind things)

Vital to this conversation of etiology is a familiarity with the complexity contributed by human variety. Social, mental, and natural elements collaborate alongside hereditary helplessness to impact an individual's general gamble for creating liquor issues. Hereditary and other organic elements are possible associated with the development of liquor reliance, alongside mental, conduct, demeanour, mental, and socio social variables.

The normal example of week after week or month to month compensation being circulated in real money prompts people with affinity for liquor maltreatment to spend huge measures of the prepared money accessible on the day, prompting shortage of cash for garments, food, schooling of kids, wellbeing and other fundamental family needs. Frequently individuals need to get cash at high paces of revenue and can be crashed into destitution. These hitting the bottle hard episodes can likewise prompt abusive behaviour at home, street car crashes, non-attendance and other such issues. As per Benegal V. (2009) liquor subordinate People spent more than they procured. A great many people take credits to help their propensity, working days and some lose their positions. Such families were upheld by pay from other relatives - and they sent youngsters under 15 to attempt to enhance family pay.

Liquor forces a high monetary expense on society. One gauge puts the yearly monetary expense of liquor maltreatment in the US to be US $ 148 billion, including US$ 19 billion for medical care use. In Canada, the financial expense of liquor adds up to roughly US$ 18.4 billion, addressing 2.7 percent of the GDP. Concentrates on in different nations have assessed the expense of liquor related issues to associate with one percent of the GDP. One of the damages connected with liquor is the unfriendly effect on the monetary status of the group of the liquor victimizer. There is a rising degree of consumption by the liquor junkie to support his/her propensity. Bit by bit, because of the limitations that the family pay forces, the liquor fiend starts to get cash, take as well as sell family objects to support his/her propensity. Frequently the passing of a task because of poor word related execution further influences the family pay unfavourably.

Flensborg-Madsen, T., Knop, J., Mortensen, E. L., Becker, U., & Grønbæk, M. (2007). Amount of alcohol consumption and risk of developing alcoholism in men and women. *Alcohol & Alcoholism*-mentioned that After change for all putative confounders, the gamble of liquor abuse for ladies expanded fundamentally at 1-7 beverages each week with a HAZARD RATIO (HR) of 2.02 (95% certainty stretch (CI): 1.16, 3.53) contrasted with never/never drinking; the HR for drinking month to month was 1.75 (95% CI: 1.08, 2.85). The gamble for men didn't increment essentially before 22-41 beverages each week (HR = 3.81, 95 % CI: 2.18, 6.68) or on the other hand on the off chance that they had an everyday liquor consumption (HR = 3.55, 95 % CI: 2.11, 5.99). Smoking was freely connected with the gamble of liquor addiction for all kinds of people. The gamble of creating liquor addiction expanded fundamentally by extremely low admissions of liquor in ladies, while the gamble is just expanded essentially in men polishing off in excess of 21 beverages each week.

Ganry, O., Baudoin, C., & Fardellone, P. (2000). Effect of alcohol intake on bone mineral density in elderly women: the EPIDOS study. *American journal of epidemiology*, *151*(8), 773-780. Liquor ingestion led to a 3-crease expansion in flowing estradiol in ladies on estrogen substitution treatment (ERT); in any case, liquor didn't change estradiol essentially in charge ladies who were not on ERT. In ladies utilizing ERT, estradiol levels expanded from 297 to 973 pmol/L (81 to 265 pg/mL) in the span of 50 minutes (P<.001) during the rising appendage of the blood liquor bend and remained fundamentally above gauge for 5 hours (P<.001). No critical expansion in circling estrone was identified in one or the other gathering. Notwithstanding, estrone levels diminished after liquor and fake treatment in ladies on ERT (P<.05). Blood liquor levels didn't contrast altogether in ladies who utilized ERT and the people who didn't. Top blood liquor levels of 21 mmol/L were accomplished in every one of the 2 gatherings

inside 50 to an hour in the wake of drinking started

Gupta, A., Shridhar, K., & Dhillon, P. K. (2015). showed that A sum of 7066 ladies matured 15-70 years showed differed degrees of mindfulness on risk factors like family ancestry (13-58%), regenerative history (1-88%) and weight (11-51%). Proficiency levels on risk factors didn't further develop over the 8-year time frame (2005-2013). On normal, medical caretakers announced higher, however still changed, mindfulness levels for risk factors like family ancestry (40.8-98%), conceptive history (21-90%) and corpulence (34-6%). Mindfulness levels were not predictably higher for the more grounded determinants of hazard. survey uncovered low disease proficiency of bosom malignant growth risk factors among Indian ladies, regardless of their financial and instructive foundation.

Gupta, P.C., Saxena, S., Pednekar, M.S., Maulik, P.K., 2003. stated Reviews of a few little locale in the upper east district of India in 1996 by Mohan what's more, partners (WHO, 2000) uncovered sex-and age-explicit paces of liquor utilize that were like the assessments revealed in the current review. They saw that as 85% of respondents detailing 'any' liquor use in the previous month were distinguished as weighty consumers who frequently announced (around 75%) having issues controlling their utilization. A comparative example was accounted for locally review of men 45 years and more established in Mumbai (Bombay), India

C. Social and Legitimate Viewpoints

In 12 significant clinics of Bangalore city India, uncovered that almost 28% of traffic wounds were straightforwardly owing to liquor. Among drivers that were being tried for liquor utilization and driving, 35% were over the lawfully admissible constraints of 30 mg/100 ml when kept an eye on breath analyser. It was the youthful male (25-39 years), proficient, with weighty drinking in bars or at parties, either alone or with companions, educated about perils of drinking however oblivious to risks or lawful outcomes, who were presenting more serious risks out and about. (NIMHANS, 2008) Statistics throughout recent many years have constrained individuals in the West to answer by development of self-improvement or care groups, for example, 'Mothers against Drunken Driving (MADD)'.

D. The family

The connection between a liquor junkie and his/her family is complicated. Relatives report encountering culpability, disgrace, outrage, dread, pain and isolation because of the presence of a liquor junkie in the family. They are frequently exposed to direct to extreme types of provocation, struggle and tense climate when they go up against the drinking conduct of their liquor manhandling relative. Life partners in families where there is ongoing, unnecessary utilization of liquor are every now and again isolated. Offspring of liquor manhandling people report a higher occurrence of close to home and school-related issues.

E. Time of Starting Drinking

Liquor addiction has generally been viewed as an issue of centre life however late examinations are recommending that liquor issue may be expanding among youngsters.As per Award, B.F., Paces of lifetime reliance declined to roughly 10% among the people who started drinking at age 20 or more seasoned. The yearly pace of this downfall was comparable for the two sexes. Albeit in the past ladies by and large began drinking at later ages than men, later review information show that this distinction has almost vanished.

Ladies' drinking is generally normal between ages 26 and 34 and among ladies who are separated or isolated. Hitting the bottle hard (i.e., utilization of at least five beverages for each event on 5 or on the other hand more days in the previous month) is generally normal among ladies ages 18 to 25 (Su, S.S.; Larison, C.; Ghadialy, R.; et al., 1997). Among racial gatherings, ladies' drinking is more common among whites, albeit people of colour are bound to drink vigorously NIAAA. (1998).

Cahalan and Casin found that men matured 21 to 41 yrs. had the most elevated rates for any age gathering of high drinking, loss of control and liquor related issues with cash, companions, neighbours and police.Robins demonstrated that non drunkards' dark guys revealed heaviest drinking at roughly age 25. Mulford and Mill operator found that the most noteworthy paces of weighty drinking in men happened in the most youthful gathering concentrated on 21 to 25 years of age, while Bailey Et Al. detailed a pace of liquor abuse for men age between 20 to 24 yrs. of 13 for each 1000.Ladies' drinking is generally normal between ages 26 and 34 and among ladies who are separated or isolated. Hard-core boozing (i.e., utilization of at least five beverages for each event on at least 5 days in the previous month is normal among ladies ages 18 to 25 years (Su, S.S. et al.,1997). Seriousness of drinking issues seems to diminish dynamically with propelling age past youthful adulthood. (Cahalan And Room, 1974)

F. Peer Tension

Most examination overall has found that by late puberty peers are regularly the most grounded effect on private way of behaving, particularly concerning liquor and substance use (Kandel, 1980, 1985).Lo's (1995) investigation of first-year understudies at a southern college tracked down that companion standards were more grounded indicators of level of inebriation than were parental standards, with peer impact being most prominent for men.Perkins'(1985) investigation of a cross-part of students at a northeastern school found peer impacts (saw companions' drinking standard and organization enrolment) to be a lot more grounded indicators of liquor utilization than other foundation factors including religion, orientation and guardians' perspectives.

The strength of friend impact might be the way to understanding discoveries where understudies will show drinking ways of behaving once in a while that they go against concerning their own perspectives (Robinson et al., 1993). Moreover, peer standards might be of specific significance in "peer-concentrated" school settings, for instance, undergrad and private foundations where understudies need regular contact with guardians.

Orocutt (1991) found that understudies arranged to drink

intensely did as such among companions. These understudies saw the presence of companions, apparently saw to be on the same page, as consolation or standardizing support for them to follow up on their drinking inclinations.

Martin and Hoffman (1993), concentrating on students at an eastern college, found that companion impact concerning the quantity of school and non-school companions who drank was a huge indicator of individual utilization even subsequent to controlling for the singular's living climate and positive hopes related with liquor use.

Young adult drinking has been related with liquor involving peers in dominatingly white examples (e.g., Kandel, 1986; Kandel and Andrews, 1987: Sieving et al., 2000). Research demonstrates that social impacts to drink come from both the underlying determination of companions (i.e., picking companions) and the ensuing support of fellowships with uplifting feedback and different prizes (Windle, 1999).

Albeit most examination has been led with prevalently white examples, a few examinations have likewise tracked down comparative connections between juvenile drinking and social elements among minority populaces. For instance, cross-sectional examinations led with dark and Hispanic teenagers showed that companion and parental use were connected with young adult drinking (e.g., Beal et al., 2001; Stream et al., 1997; Feigelman et al., 1995; Forney et al., 1991). In this manner past exploration concerning the social impacts of family and companions gives significant data about the etiology of juvenile liquor use.Research has shown that juvenile drinking standards for companions, friends, and grown-ups will quite often be erroneous. For instance, in a concentrate by Prinstein and Wang (2005), youths' discernments of their companions 'conduct unequivocally anticipated their own way of behaving. Nonetheless, view of their companions' drinking conduct was viewed as discrepant from their companions' real way of behaving, including weighty drinking.

Additionally, teenagers who misjudge the pervasiveness of wellbeing ways of behaving, for example, tobacco or liquor use, have been viewed as at higher gamble for taking part in those ways of behaving (e.g., Botvin et al., 1992.

G. The Psychological Associates

Beckman (1978) observed that alcoholic ladies were lower on confidence combined with high neuroticism, nervousness ideation.

Aubrey C. (1999) concentrated on the issue of the alcoholic self-image and observed that the mental self portrait of a heavy drinker is more negative than that of a non-alcoholic and that female alcoholic have a worse mental self-view than male drunkards.

Beckman J.L. (1999) reports that adjacent to slow confidence sex - job disarray was seen among female drunkards. Likewise, some exploration brought up to female liquor abuse and dysmenorrhea and menopause and pinpoints a connection between episodes of weighty drinking and per feminine pressure.

Soreli G.T., (1998) found that confidence and overwhelmingly male/female and manly sex composed were seen with non-alcoholic ladies though alcoholic showed levels of low confidence, manliness and womanliness and mental manliness was the central point recognizing alcoholic from non-alcoholic ladies, as well as, preferred indicators of confidence over drunkards.

Female heavy drinkers had a lower mental self-portrait than male drunkards however not in each perspective. The alcoholic ladies saw themselves all the more adversely when it came to vanity, individual self and social self. They were anyway more certain than the guys in the area of character, conduct, actual self, moral self and family self. The alcoholic male was more transparently self-basic than the alcoholic female. (Piazza N.J.,1989)

3. Health Impact

A. Hereditary Impacts

Are irrefutable in men however unsure in ladies. Heath et al (1997) consider hereditary gamble factors as a significant job in deciding liquor conditions risk in ladies as well as men, except for certain socio social factors like strict hardship, a similar character.

As yet being examined is a marker alluded to as the dopamine D2 receptor, which Blum and collaborators, Kendler, K.S., and Prescott, C.A (1997), viewed as present more frequently in heavy drinkers than in nonalcoholic Prescott, C.A. (1999). in creature studies, the dopamine D2 receptor had been related with cerebrum capabilities connecting with remuneration, support, and inspiration. In any case, various specialists have been not able to copy the aftereffects of Blum's review. (Bohman, M.; Sigvardsson, S.; also, Cloninger, C.R., 1981; Cadoret, R.J.; Yates, W.R.; Troughton, E.; et al., 1986.)

B. Physiological Impacts

Ladies retain and process liquor uniquely in contrast to men. As a rule, ladies have less body water than men of comparable body weight, so ladies accomplish higher centralizations of liquor in the blood subsequent to drinking identical measures of liquor (Frezza, M.; Di Padova, C.; Pozzato, G.; et al.1990; Taylor, J.L.; Dolhert, N.; Friedman, L.; et al. 1996).

Likewise, ladies seem to take out liquor from the blood quicker than men. This finding might be made sense of by ladies' higher liver volume per unit lean weight (Li, T.K.; Facial hair, J.D.; Orr, W.E.; et al. 1998.; Kwo, P.Y.; Ramchandani, V.A.; O'Connor, S.; et al. 1998), on the grounds that liquor is utilized nearly altogether in the liver (Levitt, M.D.; Li, R.; DeMaster, E.G.; et al., 1997). Contrasted and men, ladies foster liquor initiated liver illness over a more limited timeframe also, subsequent to polishing off less liquor (Tuyns, A.J., and Pequignot, G 1984.; Gavaler, J.S., and Arria, A.M., 1995). Also, ladies are more probable than men to foster alcoholic hepatitis and to kick the bucket from cirrhosis (Corridor, P.M., 1995). Creature research recommends that ladies' expanded gamble for liver harm might be connected to physiological impacts of the female regenerative chemical estrogen. (Ikejima, K.; Enomoto, N.; Iimuro, Y.; et al., 1998)

Breast Cancer- Many investigations report that moderate to

weighty liquor utilization expands the gamble for bosom malignant growth (Smith-Warner, S.A.; Spiegelman, D.; Yaun, S.S.; et al.1998), albeit one ongoing review found no expanded bosom disease risk related with utilization of dependent upon one beverage each day, the greatest drinking level revealed by most ladies. (Zhang, Y.; Kreger, B.E.; Dorgan, J.F.; et al 1999

4. Research Methodology

A. Need for the study

Ongoing investigations including more female subjects uncover that drinking varies among genders. Ladies acquire higher centralizations of liquor in the blood and become more weakened than men in the wake of drinking comparable measures of liquor. These have physiological results; ladies are at a higher risk for organ harm and failure than men. Furthermore, ladies' liquor reliance is viewed as less but still it harms internally. Such Distinctions in gender orientation in liquor use have reinforced expensive in how social orders recognize and attempt to control liquor related issues. This research subsequently endeavours to comprehend in a culture where ladies approach liquor, and their loves is increasing but are they really aware about the health impact of liquor.

The accompanying research questions are being tended to-

1. Which age group tend to drink the most alcohol?
2. Which occupation of females drinks the most?
3. What are Factors that influenced the increase of consumption?
4. How many women face societal judgement and criticism?
5. Do people judge the drinking habit of different women?
6. Does women know that alcohol effect differently on Men and Women
7. How many women know about the ancient alcohol consumption by women?
8. How often do women consume alcohol?

B. Objectives

1. To find whether there is an influence of entertainment industry on women's consumption of alcohol.
2. To determine the cause of alcohol intake.

C. Hypothesis

Ha_1

There is a significant number of women aged between 18-24 that consume most alcohol.

Ha_2

There is a significant number of women from IT sector that have the most Capacity

Ha_3

There is a significant number of women who prefer beer as their alcohol beverage

Ha_4

Women feel that there is societal judgement as they consume alcohol

Ha_5

There is a significant increase in alcohol consumption among women because of the influence of films, media and advertising.

D. Method

In this research investigation, while developing research design

Dependent variables are –

1. Occupation
2. Age
3. Period of consuming alcohol(period)
4. Type of alcohol beverage consumed

Independent variable

1. Level of alcohol consumption/ capacity

1) Population and Sample

This research investigates about females on various factors influencing alcohol consumption and relationship among occupation, age, period of consuming alcohol and type of alcohol beverage consumed.

The sample selected was 100 females of India and it will be a non-probability sampling.

Tools and measures the questionnaire was prepared with the help of google forms and total 100 responses were taken. Spss and excel was used for data analysis.

2) Nature of Data

Primary data- was collected with the help a survey made on Google form with 20+ questions

Secondary data- was collected from the Google scholar and research gate , articles , library , blogs , Shodhgana, thesis.

5. Results and Analysis of Data

A. Results of Data

The aim of the study is to devote research attention to women's drinking factors, about judgement, their knowledge about the ancient women consuming alcohol, health problems they are aware of, and if they feel that there is a gender specific difference in consuming alcohol.

1) Respondent Profile



Fig. 1.

According to the fig. 1, the age category of "18-24" are (37%) of the whole respondents, and the age category of "25-34" are (26%) of the whole respondents, the age category of "35-44" are (18%) of the whole respondents, the age category of "44+" are (19%) of the whole respondents.

Table 1
Your origin state?

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Rajasthan	28	28.0	28.0	28.0
	Uttar Pradesh	20	20.0	20.0	48.0
	Punjab	6	6.0	6.0	54.0
	Haryana	2	2.0	2.0	56.0
	Maharashtra	11	11.0	11.0	67.0
	Delhi	9	9.0	9.0	76.0
	Chandigarh	2	2.0	2.0	78.0
	Gujarat	4	4.0	4.0	82.0
	Uttarakhand	4	4.0	4.0	86.0
	West Bengal	4	4.0	4.0	90.0
	Assam	4	4.0	4.0	94.0
	Bihar	2	2.0	2.0	96.0
	Madhya Pradesh	2	2.0	2.0	98.0
	Jharkhand	1	1.0	1.0	99.0
	Andhra Pradesh	1	1.0	1.0	100.0
	Total	100	100.0	100.0	



Fig. 2.

According to the fig. 2 the students are 25% of the total respondents, IT Sectors are 25%, Homemaker are 19%, Industrial Professional are 18% and other are 13% of the total respondents.

Table 1 shows that 28 out of 100 respondents were from Rajasthan and 20 were from Uttar Pradesh. 11 respondents were from Maharashtra. Only 1 respondent were from Jharkhand and Andhra Pradesh.



Fig. 3.

Fig. 3 indicates that out of 100 respondents 66% women don't have any knowledge about the ancient women consuming and 34% have the knowledge.



Fig. 4.

The above Fig. 4 indicates that out of 100 women 3% women consume alcohol daily. 26% consume rarely, and most of the women consume (30%) occasionally with 22% women weekly.



Fig. 5.

The above Fig. 5 indicates that out of 100 women most women that is 33% of them prefer beer and wine 17% followed by vodka 16%, 11% Rum and cocktail. Women prefer other drinks are 13%.



Fig. 6.

Above Fig. 6 indicates that out of 100 women 39% women consume 1 drink in one siting, 2-3 drinks by 47% women and 14% women consume 4+ drinks.



Fig. 7.

Fig. 7 shows that 41% of the women were consuming alcohol from more than a year, 23% few months, 19% one year and 17% recently.



Fig. 8.

Fig. 8 shows that most women prefer 53 and 52 marked the factor as fun and social gathering, while 37 women because of work life and 29 women due to stress.



Fig. 9.

The above Fig. 9 indicates that 46% of the women think that alcohol impact differently on men & women and 33% of them are not sure while 21% says no.



Fig. 10.

Fig. 10 shows that liver disease is the most common health problem that women are aware of, while weight loss and weight gain is known by the least respondents by 29 and 30. Sleep disturbance is known by 55 women



Fig. 11.

The above Fig. 11 indicates the out of 100 respondents 72% of the them are not affected in relationship in work or other aspects of life due to their drinking habits.



Fig. 12.

The above Fig. 12 shows that 41% of the women face societal judgement or criticism whereas 37% of them responded maybe and 22% no.



Fig. 13.

Fig. 13 indicates that 51% of the women are judged because of drinking habits and 13% said no with 36% maybe.



Fig. 14.

The above Fig. 14 shows that 61% women agree that women are consuming more alcohol now a days and 12% responded no while 27% are not sure.



Fig. 15.

Fig. 15 indicates that 64% of the women out of 100 responded that films, media influences women to consume alcohol while 24% are not sure and 12% said no.



Fig. 16.

Fig. 16 indicates that 21% of the women have personally experienced health problems and 48% responded no while 31% are not sure.



Fig. 17.

Fig. 17 indicates that 73% of the women does not feel any challenge in accessing healthcare or treatment while 27% said Yes



Fig. 18.

Fig. 18 out of 100 women 93 of them responded no when asked that if they are seeking any help for health issue and 6 of them said yes



Fig. 19.

Fig. 19 indicates that 47 women feels that their family and friends are supportive & concerned about their alcohol consumption



Fig. 20.

Fig. 20 indicated that 47% women feel that there are challenges that women face when seeking healthcare or support in society while 28% said maybe



Fig. 21.

Fig. 21 indicated that 43% women strongly agree, 2%strongly disagree,1% disagree.

6. Findings & Conclusions

Ha_1-There is a significant number of women aged between 18-24 that consume most alcohol.

Table 2

Crosstab

Count

		AGE				Total
		18-24	25-34	35-44	44+	
How many drinks do you usually drink while sitting?	1 DRINK	18	13	5	3	39
	2-3 DRINKS	14	11	10	12	47
	4+ DRINKS	5	2	3	4	14
Total		37	26	18	19	100

From the above the Table 2 it is indicated that the women aged between 18-24 are consuming the most alcohol and there drinking capacity is also maximum that is more than 4 drinks.

Table 3

How often do you consume alcohol??? * AGE Crosstabulation

Count

		AGE				Total
		18-24	25-34	35-44	44+	
How often do you consume alcohol???	DAILY	1	1	0	1	3
	WEEKLY	5	7	5	5	22
	MONTHLY	5	5	4	5	19
	OCASSIONALY	14	6	5	5	30
	RARELY	12	7	4	3	26
Total		37	26	18	19	100

From the above Table 3 indicates that the women aged between 25-34 consume age the most weekly and occasionally by 18-24.

Thus, fail to reject Ha_1.

Ha_2-There is a significant number of women from IT sector that have the most Alcohol Intake Capacity.

Table 4

Crosstab

Count

		OCCUPATION					
		STUDENT	IT SECTOR	HOMEMAKER	INDUSTRIAL PROFESSIONAL	OTHER	Total
How many drinks do you usually drink while sitting?	1 DRINK	13	12	4	8	2	39
	2-3 DRINKS	10	9	12	7	9	47
	4+ DRINKS	2	4	3	3	2	14
Total		25	25	19	18	13	100

From the Table 4 it is clear that the no. of students and it sectors women are same but the intake of alcohol is most in the IT sector as they consume 4+ drinks.

Thus, fail to reject Ha_2.

Ha_3-There is a significant number of women who prefer beer as their preference in the alcohol beverage compared with occupation.

Table 5

What type of alcohol beverage do you usually drink? * OCCUPATION Crosstabulation

Count

		OCCUPATION					
		STUDENT	IT SECTOR	HOMEMAKER	INDUSTRIAL PROFESSIONAL	OTHER	Total
What type of alcohol beverage do you usually drink?	BEER	13	8	5	2	4	32
	WINE	1	5	2	9	0	17
	VODKA	6	5	1	1	3	16
	RUM	1	3	5	1	1	11
	COCKTAIL	4	4	6	5	5	24
Total		25	25	19	18	13	100

Table 5 indicates beer is preferred by 37 women and students prefer the most beer.

Thus, fail to reject Ha_3.

Ha_4 - Women feel that there is societal judgement as they consume alcohol.

From fig. 12, it is clearly seen that 41 women feel societal Judgement & criticism after seeing their alcohol habit and 37 are not sure.

Thus, fail to reject Ha_4

Ha_5 - There is a significant increase in alcohol consumption among women because of the influence of films, media and advertising.

Fig. 15 indicates that women think there is an influence of films, media & advertising on alcohol consumption.

Thus, fail to reject Ha_5.

7. Conclusion

1) This research objective was to find whether there is an influence of film, media and advertising on women alcohol consumption total 100 women were given the survey out of which 64 women said yes that there is an influence of alcohol consumption because of films, media and advertising.
2) There are different reason because of which women consume alcohol like 1 = Fun 2=Life events 3=work life 4=stress 5=emotional 6=social pressure 7=social gathering. From the research it was shown that all the factors influence the consumption of alcohol but the most common factor was Fun and social gathering
3) Most of alcohol is consumed by the age group of 18-24 years. Women aged from 18-24 drink maximum for fun
4) Students (25) & IT Sectors (25) women consumes the most alcohol. Conclusion from this is that students & IT sector are more prone to the health effect that may be caused because of alcohol consumption
5) Factor that have influenced the alcohol consumption the most is the Fun and Social Gathering, stress, social pressure are also some common factors that are seen in then women.
6) There are total 41 women who face societal judgement and 37 of them are not sure about it 22 women said that they do not face any societal judgement at all. The conclusion drawn is that maximum women face societal judgement
7) 51 women agrees that people judge them when they share their drinking habits with other people and 36 of them are not sure and 13 said no So, from this the conclusion is that women face judgement because of drinking habits
8) 56 women feel that alcohol impact them differently than men and 33 selected maybe 21 women said no so from this it is clear that many women know that alcohol react differently to different genders
9) Only 34 women know about the ancient alcohol consumption by women and 66 of them does not have any knowledge so from this the conclusion is that women do not have any knowledge about the past history of Indian alcohol consumption by women
10) Maximum women tend to drink alcohol occasional and some of them weekly

To briefly conclude about this research there are women who does not have much awareness about the health problems but still they tend to consume more alcohol and also women do not have any knowledge about the ancient alcohol consumption by women and today also women face societal judgement because of their drinking habits. The increase in consumption is seen in the age group of 18-24 and among students and IT sector. Most women are aware about the health effects. The factor to consume alcohol tend to be many but fun and social gathering is the most common.

8. Recommendations

1) There should not be any societal judgement on women because of her drinking habits as because of this she may found it difficult to consult for any health problems
2) Family and friends should support the women if they are seeking any healthcare
3) Ancient knowledge about alcohol consumption should be there
4) A proper knowledge should be given on the impact of alcohol on women by the government
5) Entertainment industry should not show content that may influence the women to consume alcohol.

9. Implications

A. Government

1) This research is of great use to the government as they can improve their health care service and support for women consuming alcohol.

2) Government can use this research and aware people about the health problems caused by alcohol consumption

B. Society

1) They can use this research to understand the pattern of alcohol consumption among women.

C. Public Health

1) This research can help the inform targeted prevention and intervention strategies to mitigate these risks. Increased alcohol sensitivity in women can lead to faster and more severe alcohol-related health problems like liver damage, heart disease, and certain cancers.
2) Women experiencing stress, trauma, or mental health challenges may be particularly susceptible to increased alcohol sensitivity. Research can help identify and support these vulnerable groups**.**

D. Individual Well-being

1) Personalized treatment: Understanding the factors behind increased alcohol sensitivity can pave the way for personalized treatment approaches for women struggling with alcohol misuse. This can lead to more effective and lasting recovery outcomes.

Empowerment and education: Research findings can empower women to make informed choices about alcohol consumption and seek help if needed. Educational campaigns can raise awareness about the specific risks women face and encourage responsible drinking practices.

10. Limitations

Despite detail study and careful implementation of research design, there are many limitations to the study. Most of the limitations occurred because of the nature of the topic that was investigating.

Present below some of the limitation in the dissertation:

1) Additionally to discuss this behaviour many women find it difficult to share because of the taboo associated with drunkenness
2) The study was conducted on a convenient sample
3) The sample was small because of time constrain

11. Scope for the Future Research

1) The above research study can be used on a larger sample of women with particularly on one state.
2) The connection of stress and alcohol consumption can be explored further.

Women alcoholics have historically been underrepresented in research studies certainly not sufficiently in proportion to the differences in male/female. This suggests the subtle sex bias that exists while dealing with women who have drinking problems.

Thus, in conclusion bringing women in as research subject in alcohol studies is a step towards addressing the emerging concerns of women with drinking problems in India.

References

[1] E. L. Abel and M. Kruger, "Physician attitudes concerning legal coercion of pregnant alcohol and drug abusers," *American Journal of Obstetrics and Gynecology*, vol. 186, pp. 768–772, 2002.

[2] M. Alam and A. Karan, *Elderly Health in India: Dimension, Differentials and Determinants*. New Delhi, India: United Nations Population Fund (UNPA), 2011.

[3] N. Almeida-Filho, I. Lessa, L. Magalhães, M. J. Araújo, E. Aquino, I. Kawachi, and S. A. James, "Alcohol drinking patterns by gender, ethnicity, and social class in Bahia, Brazil," *Revista de Saúde Pública*, vol. 38, pp. 45–54, 2004.

[4] C. Aubrey, G. Mc, L. Romo, and S. Joffre, *The Alcoholic Self Image: A Comparative Study Between Men and Women*. Paris, France: University Paris VIII, 1999.

[5] Australian Institute of Health and Welfare, *1998 National Drug Strategy General Household Survey: First Results*. Canberra, ACT, Australia: Department of Health and Aged Care, 1999.

[6] Baiely *et al*., cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.

[7] V. Benegal, G. Gururaj, and P. Murthy, Report on a WHO Collaborative Project on Unrecorded Consumption of Alcohol in Karnataka, India. World Health Organization, 2003.

[8] S. B. Blume, "Women and alcohol: Issues in social policy," in *Gender and Alcohol: Individual and Social Perspectives*, R. W. Wilsnack and S. C. Wilsnack, Eds. New Brunswick, NJ, USA: Rutgers Center of Alcohol Studies, 1997, pp. 462–489.

[9] M. Bohman, S. Sigvardsson, and C. R. Cloninger, "Maternal inheritance of alcohol abuse: Cross-fostering analysis of adopted women," *Archives of General Psychiatry*, vol. 38, no. 9, pp. 965–969, 1981.

[10] R. S. Brienza and M. D. Stein, "Alcohol use disorders in primary care: Do gender-specific differences exist?" *Journal of General Internal Medicine*, vol. 17, pp. 387–397, 2002.

[11] Cahalan and Casin, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.

[12] Cahalan and Room, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.

[13] Centers for Disease Control and Prevention, Alcohol and Public Health: Alcohol-Related Disease Impact—Annual Average for United States 2015–2019 Alcohol-Attributable Deaths Due to Excessive Alcohol Use. Atlanta, GA, USA: CDC, 2022.

[14] Y. M. Cha, "Women more vulnerable to negative effects of alcohol," *Alcoholism: Clinical and Experimental Research*, Jan. 2006.

[15] H. P. Chalfant and B. S. Roper, Social and Behavioral Aspects of Female Alcoholism: An Annotated Bibliography. Westport, CT, USA: Greenwood Press, 1980.

[16] I. Chein, "Psychological functions in drug use," in *Scientific Basis of Drug Dependence*, Steinberg, Ed. London, U.K.: Churchill Ltd., 1969.

[17] Clark, cited in C. R. Eng and D. Hanson, "Gender differences in drinking patterns and problems among college students: A review of the literature," *Journal of Alcohol and Drug Education*, vol. 35, no. 2, pp. 36–47, 1990.

[18] W. Doniger, *The Laws of Manu*. New Delhi, India: Penguin Classics, 1992.

[19] S. Edith and L. Gomberg, "Women and alcohol: Use and abuse," *Journal of Nervous and Mental Disease*, vol. 181, no. 4, pp. 211–219, Apr. 1993.

[20] E. E. Epstein, K. Fischer-Elber, and Z. Al-Otaiba, "Women, aging, and alcohol use disorders," *Journal of Women & Aging*, vol. 19, no. 1–2, pp. 31–48, 2007.

[21] A. Erol and V. M. Karpyak, "Sex and gender-related differences in alcohol use and its consequences: Contemporary knowledge and future research considerations," *Drug and Alcohol Dependence*, vol. 156, pp. 1–13, 2015.

[22] T. Fairley, "The early history of distillation," *Journal of the Institute of Brewing*, vol. 13, no. 6, pp. 559–582, 1907.

[23] P. Fergusson, T. Lemaron, *et al*., *Drugs and Family/Peer Influence on Adolescent Drug Use*. Rockville, MD, USA: National Institute on Drug Abuse, 1975.

[24] K. M. Fillmore, J. M. Golding, E. V. Leino, M. Motoyoshi, C. Shoemaker, H. Terry, *et al*., "Patterns and trends in women's and men's drinking," in *Recent Developments in Alcoholism*, 1997.

[25] M. Frezza, C. Di Padova, G. Pozzato, *et al*., "High blood alcohol levels in women: The role of decreased gastric alcohol dehydrogenase activity and

first-pass metabolism," *The New England Journal of Medicine*, vol. 322, no. 2, pp. 95–99, 1990.
[26] From, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.
[27] Goodwin and G. Gabrielli, cited in G. Miller, *Learning the Language of Addiction Counseling*. Hoboken, NJ, USA: John Wiley & Sons, 2000.
[28] P. M. Hall, "Factors influencing individual susceptibility to alcoholic liver disease," in *Alcoholic Liver Disease: Pathology and Pathogenesis*, 2nd ed., P. M. Hall, Ed. London, U.K.: Edward Arnold, 1995, pp. 299–316.
[29] D. J. Hanson, "History of alcohol and drinking around the world." [Online]. Available: http://www.potsdam.edu/hansondj
[30] A. C. Heath, K. K. Bucholz, P. A. Madden, S. Dinwiddie, W. S. Slutske, L. J. Bierut, *et al*., "Genetic and environmental contributions to alcohol dependence risk in a national twin sample: Consistency of findings in women and men," *Psychological Medicine*, vol. 27, pp. 1381–1396, 1997.
[31] R. Hingson, W. Zha, B. Simons-Morton, and A. White, "Alcohol-induced blackouts as predictors of other drinking-related harms among emerging young adults," *Alcoholism: Clinical and Experimental Research*, vol. 40, no. 4, pp. 776–784, 2016.
[32] E. L. Hornick, *The Drinking Woman*. New York, NY, USA: Association Press, 1977.
[33] K. Ikejima, N. Enomoto, Y. Iimuro, *et al*., "Estrogen increases sensitivity of Kupffer cells to endotoxin," *Alcoholism: Clinical and Experimental Research*, vol. 22, no. 3, pp. 768–769, 1998.
[34] India Finance Pvt. Ltd., "India Finance," Apr. 2002, pp. 1–3.
[35] H. Kalant, "Absorption, diffusion, distribution and elimination of ethanol: Effects on biological membranes," in *The Biology of Alcoholism*, vol. 1, B. Kissin and H. Begleiter, Eds. New York, NY, USA: Plenum Press, 1971.
[36] R. Kandel and Y. Lo, cited in H. W. Perkins, "Social norms and the prevention of alcohol misuse in collegiate contexts," *Journal of Studies on Alcohol*, Supplement no. 14, 2002.
[37] K. S. Kendler and C. A. Prescott, "Population-based twin study of alcohol abuse and dependence: Modeling gender differences," *American Journal of Medical Genetics*, vol. 74, no. 6, p. 574, 1997.
[38] G. Knupfer, "Use of alcoholic beverages by society and its cultural implications," *California's Health*, vol. 18, pp. 17–21, 1960.
[39] V. S. Kumar, "Behavioural malignancy: Alcoholism, a bleak future?" in *Bioethics in India: Proceedings of the International Bioethics Workshop in Madras*, J. Azariah, H. Azariah, and D. R. J. Macer, Eds. Madras, India: Eubios Ethics Institute, 1997.
[40] M. D. Levitt, R. Li, E. G. DeMaster, *et al*., "Use of measurements of ethanol absorption from stomach and intestine to assess human ethanol metabolism," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 273, no. 4, pp. G951–G957, 1997.
[41] T. K. Li, J. D. Beard, W. E. Orr, *et al*., "Gender and ethnic differences in alcohol metabolism," *Alcoholism: Clinical and Experimental Research*, vol. 22, no. 3, pp. 771–772, 1998.
[42] M. Jaganmohan, "Production volume of ethanol in India 2011–2020," *Statista*, 2022.
[43] A. Malhotra and A. Mohan, "National policies to meet the challenge of substance abuse: Programmes and implementation," *Indian Journal of Psychiatry*, vol. 42, pp. 370–377, 2000.
[44] H. Malin, J. Coakley, and C. Kaelber, "An epidemiologic perspective on alcohol use and abuse in the United States," in *Alcohol Consumption and Related Problems*, National Institute on Alcohol Abuse and Alcoholism, Alcohol and Health Monograph No. 1, DHHS Publ. No. (ADM) 82-1190. Washington, DC, USA: U.S. Government Printing Office, 1982, pp. 99–153.
[45] P. Martin and M. Hoffman, "Peer influence and alcohol consumption among university undergraduates," unpublished study, 1993.
[46] McCord and O. H. Mowrer, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.
[47] D. Mohen *et al*., *Current Research in Drug Abuse in India*. New Delhi, India: Proceedings of the 2nd National Conference, 1980.
[48] K. J. Mukamal, J. S. Tolstrup, J. Friberg, G. Jensen, and M. Grønbæk, "Alcohol consumption and risk of atrial fibrillation in men and women: The Copenhagen City Heart Study," *Circulation*, vol. 112, no. 12, pp. 1736–1742, 2005.
[49] Mulford and Miller, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.
[50] P. Murthy, N. Manjunatha, B. N. Subodh, P. K. Chand, and V. Benegal, "Substance use and addiction research in India," *Indian Journal of Psychiatry*, vol. 52, Suppl. 1, pp. S189–S199, 2010.
[51] R. J. M. Neve, P. H. Lemmens, and M. J. Drop, "Gender differences in alcohol use and alcohol problems: Mediation by social roles and gender-role attitudes," *Substance Use & Misuse*, vol. 32, pp. 1439–1459, 1997.
[52] S. Naik, *Indian Spirits Market: Regulatory Shackles Easing*. Mumbai, India: Rabobank International, 2002.
[53] National Institute on Alcohol Abuse and Alcoholism, *Drinking in the United States: Main Findings from the 1992 National Longitudinal Alcohol Epidemiologic Survey*, vol. 6. Rockville, MD, USA: NIAAA, 1998.
[54] National Institute on Alcohol Abuse and Alcoholism, *Alcohol Alert No. 45: Alcohol and Coronary Heart Disease*. Rockville, MD, USA: NIAAA, 1999. [Online]. Available: http://www.niaaa.nih.gov
[55] National Institute on Alcohol Abuse and Alcoholism, *Alcohol Alert No. 63: Alcohol Damaging Effects on the Brain*. Rockville, MD, USA: NIAAA, 2004. [Online]. Available: http://www.niaaa.nih.gov
[56] Orcutt, cited in H. W. Perkins, "Social norms and the prevention of alcohol misuse in collegiate contexts," *Journal of Studies on Alcohol*, Supplement no. 14, 2002.
[57] N. J. Piazza, J. L. Vrbka, and R. D. Yeager, "Telescoping of alcoholism in women alcoholics," *International Journal of the Addictions*, vol. 24, no. 1, pp. 19–28, 1989.
[58] S. Ravindran, "The Swinging Janes," *Outlook Express*, 2010.
[59] R. Ray and H. K. Sharma, "Drug addiction: An Indian perspective," in *Souvenir of ANCIPS 1994*, V. P. Bashyam, Ed. Madras, India: Indian Psychiatric Society, 1994, pp. 106–109.
[60] S. Ray, "Why women get addicted to alcohol," *The Tribune*, Apr. 21, 2002.
[61] L. N. Robins and L. T. McEvoy, "Conduct problems as predictors of substance abuse," in *Straight and Devious Pathways from Childhood to Adulthood*, L. N. Robins and M. Rutter, Eds. Cambridge, U.K.: Cambridge University Press, 1990.
[62] Robinson *et al*., cited in H. W. Perkins, "Social norms and the prevention of alcohol misuse in collegiate contexts," *Journal of Studies on Alcohol*, Supplement no. 14, 2002.
[63] R. Room, J. A. Babor, and J. Rehm, "Alcohol and public health," *The Lancet*, vol. 365, no. 9458, pp. 519–530, 2005.
[64] R. Room, A. Janca, L. A. Bennett, *et al*., "WHO cross-cultural applicability research on diagnosis and assessment of substance use disorders: An overview of methods and selected results," *Addiction*, vol. 91, pp. 199–220, 1996.
[65] Rosenberg, I. Chein, and S. B. Blume, cited in J. Chunkapura, *Alcoholism—Psychosocial Variables*, Ph.D. dissertation, University of Delhi, Delhi, India, Archana Printers, Kottayam, India, 1998.
[66] M. Sandmaier, The Invisible Alcoholics: Women and Alcohol Use in America. New York, NY, USA: McGraw-Hill, 1980.
[67] N. L. Singh, P. K. Mishra, S. K. Shukla, J. Kumar, and R. Singh, "Alcoholic fermentation techniques in early Indian tradition," 2010.
[68] S. A. Smith-Warner, D. Spiegelman, S. S. Yaun, *et al*., "Alcohol and breast cancer in women: A pooled analysis of cohort studies," *JAMA*, vol. 279, no. 7, pp. 535–540, 1998.
[69] G. T. Soreli, L. Y. Silvia, and N. A. Busch-Rossnagel, "Sex-role orientation and self-esteem in alcoholic and non-alcoholic women," Department of Human Development and Family Studies, Texas Tech University, Lubbock, TX, USA, 1998.
[70] J. L. Taylor, N. Dolhert, L. Friedman, *et al*., "Alcohol elimination and simulator performance of male and female aviators: A preliminary report," *Aviation, Space, and Environmental Medicine*, vol. 67, no. 5, pp. 407–413, 1996.
[71] A. J. Tuyns and G. Pequignot, "Greater risk of ascitic cirrhosis in females in relation to alcohol consumption," *International Journal of Epidemiology*, vol. 13, no. 1, pp. 53–57, 1984.
[72] A. Urbano-Marquez, R. Estruch, J. Fernández-Solá, *et al*., "The greater risk of alcoholic cardiomyopathy and myopathy in women compared with men," *JAMA*, vol. 274, no. 2, pp. 149–154, 1995.
[73] H. Viswanathan and J. A. Wilson, "Alcohol—The neglected risk factor in head and neck cancer," *Clinical Otolaryngology & Allied Sciences*, vol. 29, no. 4, pp. 295–300, 2004.
[74] G. J. Wang, N. D. Volkow, J. S. Fowler, *et al*., "Regional cerebral metabolism in female alcoholics of moderate severity does not differ from that of controls," *Alcoholism: Clinical and Experimental Research*, vol. 22, no. 8, pp. 1850–1854, 1998.

[75] A. White, I. J. P. Castle, C. M. Chen, M. Shirley, D. Roach, and R. Hingson, "Converging patterns of alcohol use and related outcomes among females and males in the United States, 2002–2012," *Alcoholism: Clinical and Experimental Research*, vol. 39, pp. 1712–1726, 2015.

[76] R. W. Wilsnack, S. C. Wilsnack, and A. D. Klassen, "Women's drinking and drinking problems: Patterns from a 1981 national survey," *American Journal of Public Health*, vol. 74, no. 11, pp. 1231–1238, 1984.

[77] S. C. Wilsnack, "Drinking: Recent findings and some implications for prevention," in *Women and Alcohol: Issues for Prevention Research*, NIH Publication No. 96, 1996, p. 19.

[78] S. C. Wilsnack and L. J. Beckman, *Alcohol Problems in Women: Antecedents, Consequences, and Intervention*. New York, NY, USA: Guilford Press, 1984.

[79] M. Windle, "An alcohol involvement typology for adolescents: Convergent validity and longitudinal stability," *Journal of Studies on Alcohol*, vol. 57, pp. 627–637, 1996.

[80] World Health Organization, The World Health Report 2002: Reducing Risks, Promoting Healthy Life. Geneva, Switzerland: WHO, 2002.

[81] World Health Organization, Technical Brief: Assessment of Burden and Socio-Economic Cost Attributable to Alcohol Use in India. Geneva, Switzerland: World Health Organization, 2018.

[82] Y. Zhang, B. E. Kreger, J. F. Dorgan, *et al*., "Alcohol consumption and risk of breast cancer: The Framingham Study revisited," *American Journal of Epidemiology*, vol. 149, no. 2, pp. 93–101, 1999.